

Geological Field Trips

2016

Vol. 8 (2.2)



ISSN: 2038-4947



*Società Geologica
Italiana*



ISPRA

Dipartimento per il
SERVIZIO GEOLOGICO D'ITALIA
Organo Cartografico dello Stato (legge n°68 del 2-2-1969)

A trip through the Wildhorn Nappe from Cretaceous to Neogene time
(Helvetic Nappes, Switzerland)

Tectonic Studies Group of the Swiss Geological Society - Summer excursion, 2015

DOI: 10.3301/GFT.2016.04

GFT - *Geological Field Trips*

Periodico semestrale del Servizio Geologico d'Italia - ISPRA e della Società Geologica Italiana
Geol.F.Trips, Vol.8 No.2.2 (2016), 30 pp., 20 figs. (DOI 10.3301/GFT.2016.04)

A trip through the Wildhorn Nappe from Cretaceous to Neogene time (Helvetic Nappes, Switzerland)

Tectonic Studies Group of the Swiss Geological Society, Summer excursion, 8th – 9th August, 2015

Giovanni Luca Cardello*, Neil Mancktelow**

* Section des sciences de la Terre et de l'environnement, Université de Genève, Rue des Maraîchers 13, CH-1205 Genève

** Department of Earth Sciences, ETH, CH-8092 Zürich, Switzerland

Corresponding Author e-mail address: luca.cardello@unige.ch

Responsible Director
Claudio Campobasso (ISPRA-Roma)

Editor in Chief
Gloria Ciarapica (SGI-Perugia)

Editorial Responsible
Maria Letizia Pampaloni (ISPRA-Roma)

Technical Editor
Mauro Roma (ISPRA-Roma)

Editorial Manager
Maria Luisa Vátovec (ISPRA-Roma)

Convention Responsible
Anna Rosa Scalise (ISPRA-Roma)
Alessandro Zuccari (SGI-Roma)

Editorial Board

*M. Balini, G. Barrocu, C. Bartolini,
D. Bernoulli, F. Calamita, B. Capaccioni,
W. Cavazza, F.L. Chiocci,
R. Compagnoni, D. Cosentino,
S. Critelli, G.V. Dal Piaz, C. D'Ambrogi,
P. Di Stefano, C. Doglioni, E. Erba,
R. Fantoni, P. Gianolla, L. Guerrieri,
M. Mellini, S. Milli, M. Pantaloni,
V. Pascucci, L. Passeri, A. Peccerillo,
L. Pomar, P. Ronchi, B.C. Schreiber,
L. Simone, I. Spalla, L.H. Tanner,
C. Venturini, G. Zuffa.*

ISSN: 2038-4947 [online]

<http://www.isprambiente.gov.it/it/pubblicazioni/periodici-tecnici/geological-field-trips>

The Geological Survey of Italy, the Società Geologica Italiana and the Editorial group are not responsible for the ideas, opinions and contents of the guides published; the Authors of each paper are responsible for the ideas, opinions and contents published.

Il Servizio Geologico d'Italia, la Società Geologica Italiana e il Gruppo editoriale non sono responsabili delle opinioni espresse e delle affermazioni pubblicate nella guida; l'Autore/i è/sono il/i solo/i responsabile/i.

INDEX

Information

Abstract/Riassunto	4
Program/Accommodation/Useful number	6

Excursion notes

Geological background	7
------------------------------------	----------

Itinerary

First day: nappe boundaries, syn-sedimentary structures and Neogene oblique-normal faults	10
--	-----------

STOP 1 - A panoramic overview of the Ultrahelvetic basal thrust	11
STOP 2 - The Chamossaire fault	12
STOP 3 - The Sex Rouge-Donin palaeo-fault	13
STOP 4 - Half-graben structures.....	16
STOP 5 - Latest Aptian syn-sedimentary dykes	17
STOP 6 - An example of a Neogene fault	17
STOP 7 - Turonian slumps within Campanian marls	17
STOP 8 - A panoramic view of the syn-sedimentary structures	20
STOP 9 - Neogene vein sets	20

Second day: a closer look at the Cretaceous syn-sedimentary fault system	21
---	-----------

STOP 10 - The escarpment: a stratigraphic contact	21
STOP 11 - Palaeo-escarpment culmination	21
STOP 12 - Palaeo-escarpment and buttressing	23
STOP 13 - Slumps	23
STOP 14 - A way-back trip through the Cretaceous series	23
STOP 15 - Palaeo-geographic Maastrichtian evidences	25

References	28
-------------------------	-----------

Abstract

A field trip across the Wildhorn Nappe (SW Switzerland), forming part of the Helvetic nappes, is presented here. The field trip, which took place on the 8-9th August 2015, was the annual excursion of the Tectonic Studies Group of the Swiss Geological Society. It has been organized as a 2-day itinerary along the southern slopes of the Helvetic Alps between Kanton Bern and Valais (Fig. 1). This guide contains a brief structural and stratigraphic overview followed by Stop descriptions along a main geological cross-section that, from south to north, shows both the Neogene and the Cretaceous faults and their different structural characteristics. Neogene faults are associated with veins and ductile to brittle structures. Cretaceous faults are comparatively dry, discrete and related to palaeo-escarpments associated with stratigraphic unconformities, fault-growth geometries, slumps and sedimentary dykes. The temporal evolution of the Cretaceous syn-sedimentary fault system is also documented by spectacular panoramas and briefly discussed in relation to nappe-stack development and subsequent Neogene orogen-parallel extension.

Key words: Helvetic nappes, brittle-ductile faults, veins, syn-sedimentary faults, palaeo-escarpments

Riassunto

In questa guida presentiamo un percorso attraverso la Falda del Wildhorn (Svizzera sudoccidentale), una delle falde strutturalmente più alte delle coperture elvetiche. L'escursione ha preso luogo tra l'8 e il 9 agosto 2015 in quanto scelta come escursione annuale del Gruppo Studi Tettonici della Società Geologica Svizzera. Il percorso è stato organizzato come un itinerario di 2 giorni lungo i fianchi meridionali delle Alpi Elvetiche tra il Canton Berna e il Vallese (Fig. 1). Questa guida contiene anche un breve inquadramento strutturale e stratigrafico, seguito dalla descrizione degli Stop lungo una sezione geologica di riferimento che da sud a nord mostra sia le strutture del Neogene che quelle Cretacee, concentrandosi sulle loro diverse caratteristiche strutturali. Le faglie neogeniche sono associate a vene e strutture da duttili a fragili. Le faglie cretacee invece sono associate a fratture non riempite, sono discrete e relazionate a paleo-scarpate associate a contatti stratigrafici inconformi, geometrie di faglie di crescita, slumps e dicchi sedimentari. L'evoluzione temporale del sistema di faglie sin-sedimentarie cretacee è documentato anche da foto panoramiche spettacolari ed è brevemente discusso in rapporto anche allo sviluppo del sistema a pieghe e sovrascorrimenti e alla successiva estensione parallela all'asse principale dell'orogene alpino.

Parole chiave: Successioni carbonatiche, tettonica cretacea, tettonica neogenica, faglie sinsedimentarie, transizione fragile-duttile




Fig. 1 - Paths and Stops. First day path is marked by a green line and the second day by a violet line

Program

The excursion is based on public transport and walking in rather high but not difficult mountain conditions. The best time for this field trip is in the late summer when snow cover is limited. In order to have two full and efficient days, it is recommended to start already at 8:00 at the bus station of Sion Post, Place de la Gare (directly opposite the main railway station). For most participants, this will mean overnighing in Sion on the previous evening. Cheap accommodation is available at the Youth Hostel, which is very close to the train station. On reaching the cable car of Anzère, you should get out at the final station called Pas de Maimbré (Fig. 1), from where the trip continues on foot. For cable car time schedules in summer time or to book for a group write an email well in advance to info@teleanzere.ch. Overnighing in the field, dinner and breakfast are suggested at the Cabanne des Audannes, where a preliminary reservation is recommended for groups in high season. You can book at this website : <http://www.audannes.ch>.

On the afternoon of the second day, there will be a bus from Les Rousses to bring you back to Sion. Please check out all time schedules of the Swiss public transportation at <http://www.ffs.ch/home.html>. Participation in the excursion is at your own risk. The

field trip involves hiking in alpine terrain at altitudes up to ca. 2600 m. Adequate mountain gear is necessary, in particular mountain boots and warm clothing. Please take along a picnic for 2 days. More detailed topographic and geologic support is provided by the 1:25.000 St-Leonard sheet, which you may wish to borrow or buy. A pre-trip reading of Cardello & Mancktelow (2014, 2015) is recommended.

Accommodation

Auberge de Jeunesse Sion – Ostello della Gioventù di Sion. Rue de l'Industrie 2 1950 Sion (VS) – Tél: +41 27 323 74 70 – i-contacts: sion@youthhostel.ch;

Useful numbers in case of emergency

General Emergency – 112 – Emergenze generiche
 Firefighters – 118 - Pompieri
 Police – 117 - Polizia
 Ambulance – 144 - Ambulanza
 Rega (elisoccorso) – 1414 – Elisoccorso alpino



Geological background

The SW Helvetic Alps (Fig. 2) record several tectonic events such as syn-rift extension, post-break-up extension, foreland extension, orogenic compression and orogen-parallel to orogen-perpendicular extension during continued convergence. As a result, different generations of faults were produced from the Mesozoic to the Neogene. Similar

to what is observed elsewhere (e.g. Mohn et al., 2010; Cardello & Doglioni, 2015), the Helvetic succession was repeatedly affected by syn-sedimentary normal faults, some of which have considerable displacement (up to ca. 500 m) and strongly affect the distribution of sedimentary facies (e.g. Günzler-Seiffert, 1941; Hänni & Pfiffner, 2001; Kempf & Pfiffner, 2004; Cardello & Mancktelow, 2014).




Fig. 2 - **a)** Simplified tectonic map of SW Switzerland. In blue, axis of the Rawil Depression, with the plunge direction indicated (modified from Cardello 2013, after Tectonic Map of Switzerland, and Steck et al. 1999); **b)** Tectonic map of the Alps, with the location of a) indicated by the white rectangle.

During the Alpine convergence, two major stages can be distinguished (Heim, 1920; Ramsay et al., 1983; Burkhard, 1988; 1993; Jeanbourquin, 1994; Maurer et al., 1997; Pfiffner, 2009). Thin-skinned tectonics during the first stage produced folds and thrusts, with the major thrusts with displacements of (many) 10's of kilometres marking the base of distinct, named "nappes" (e.g., the Wildhorn Nappe). On a broad scale, the resultant nappe stack is composed of Penninic, Ultrahelvetetic and Helvetic nappes, which were emplaced in the time period from ca. 32 to 15 Ma (Trümpy, 1960; Crespo-Blanc et al., 1995; Steck et al., 1999; Kirschner et al., 2003). The second stage of compression is associated with cooling and with the progressive exhumation of the Alps (Burkhard, 1988; Vernon et al., 2008, Sternai et al., 2012). The nappe stack was up-domed and folded on a large scale (ca. 10 km across), with an amplitude pointing to crustal-scale thrusting (Pfiffner, 2009; Cardello et al., 2015). The axial depression of this antiform has its structurally lowest point in the Rawil pass region (i.e. the Rawil Depression; Lugeon, 1914-1918; Gasser & Mancktelow, 2010), where U-Th/He zircon age data show that cooling below 200°C occurred in the last 17-15 Ma (Cardello, 2013). The different structural style of the two major compressional events reflects a gradual change from thin- to thick-skinned tectonics within the basement, linked to ongoing deformation of the outer arc of the Western Alps (Maurer et al., 1997; Sue et al., 2007). Veins are a common feature within the Helvetic nappes (e.g. Ramsay et al., 1981; Dietrich et al., 1983; Burkhard, 1988; Cardello & Tesei, 2013; Cardello & Mancktelow, 2015) and are important markers recording the incremental stretching history (e.g. Durney & Ramsay, 1973; Casey et al., 1983). Burkhard & Kerrich (1988) distinguish three different sets of veins: pre-, syn- and post-tectonic according to their involvement in the nappe-emplacment and further deformation. However, none of the veins are strictly post-tectonic because they still reflect the incremental strain at the time of their formation.

Studies north of the Rhône Valley have mainly focused on the neotectonics and the potential relationship of faults to active seismicity. From vein relationships, Franck et al. (1984) established a succession of stretching events in the Sanetsch pass area of the Rawil Depression, relating the youngest NW-striking veins to present-day seismicity, the N-striking veins to nappe-emplacment and the NE-striking veins to syn-sedimentary faulting. During the latest stage of deformation, younger transverse faults were exhumed to the surface in the footwall of the Simplon-Rhône fault, associated with differential uplift of the Mont Blanc/Aiguilles Rouges and Aar/Gastern External Crystalline Massifs and development of the Rawil Depression. These dextral-transpressive Neogene faults generally transect pre-existing syn-sedimentary faults without significant inversion or reactivation.





Fig. 3 - Stratigraphic column of the Wildhorn Nappe, modified after Cardello & Mancktelow (2014, and references therein).



First day: nappe boundaries, syn-sedimentary structures and Neogene oblique-normal faults

The total path length of the first day is some 6.5 km, whereas the height difference is about 350 m but with some up and down. The highest point is 2766 m and the difficulty grade is medium. A few easy off-track deviations are included. From the bus station of Sion the cable car of Anzère can be reached by Post Bus. The uppermost cable car stop is Pas de Maimbré (Fig. 4a), from which you can observe the Ultrahelvetic basal thrust on the other side of the valley toward the NE (Fig. 4b, c).



From the bus station of Sion the cable car of Anzère can be reached by Post Bus. The uppermost cable car stop is Pas de Maimbré (Fig. 4a), from which you can observe the Ultrahelvetic basal thrust on the other side of the valley toward the NE (Fig. 4b, c).




Fig. 4 - **a)** Chamossaire area, the Ultrahelvetic basal thrust ramps onto the Cretaceous succession of the Wildhorn Nappe; this package is later folded together in the Prabé syncline. **b)** and **c)** Detail of the thrust zone. Asymmetric boudinage affects the rocks close to the contact, giving top-to-the-NW sense of shear.



STOP 1 - (46°18'54.03"N / 7°22'55.11"E) a panoramic overview of the Ultrahelvetic basal thrust

This Stop is a quick view of the tectonic ramp contact, which juxtaposes the Ultrahelvetic Quinten limestones over the Wildhorn Nappe (Fig. 4b, c). From here, we will walk on along the cross-section AB in Fig. 5 from the Prabé fold core to the Eaux Froides palaeo-fault.




Fig. 5 - Geological profiles (modified after Cardello & Mancktelow 2014), the trace of which are indicated in Fig. 1. Most of the faults in red that crosscut folds and nappe boundaries belong to fault set 2), as shown in Fig. 1. Palaeofaults (purple lines) determine facies and thickness variation in the Upper Cretaceous series.



STOP 2 - (46°19'5.77"N/7°22'49.88"E) the Chamossaire fault

The Chamossaire fault (Fig. 6a) is a well exposed EW-striking, steeply dipping oblique normal fault with an important dextral strike-slip component. The fault length is ca. 1.2 km and the displacement ca. 100 m. As seen from Fig. 6b, the footwall consists of: **1**) little deformed to veined limestones of the Tierwis Fm (Fig. 6b, zone a), which are crossed by sporadic, 2-3 cm thick veins of calcite immediately adjacent to the sharp contact with the fault core; **2**) fractured, veined and gradationally recrystallized limestones of the Schrattekalk fm, which are progressively dragged into the fault core, **3**) marly limestones and marls of the Wang Fm that are affected by S/C structures close to the main slip surfaces. The fault core is 1-1.5 m thick and can be divided





Fig. 6 - (modified after Cardello & Mancktelow, 2015). Chamossaire fault core (2310 m). **a**) Photograph; **b**) Interpretation: a) Fractured and veined Tierwis Fm (Lower Cretaceous marly limestones); b) Foliated and fractured cataclasite with clasts of Tierwis and Schrattekalk fm; c) Fractured and recrystallized breccia of Schrattekalk fm (Lower Cretaceous platform limestones); d) Riedel fractures and asymmetric folds in foliated cataclasite indicating dextral shear, with repeated faulting and more ductile shear; e) Foliated Wang Fm marls (Lower Maastrichtian).



into 4 zones characterized by different structure and therefore rheological behaviour during the fault activity. The folding of Riedel fractures and discrete slip surfaces within the fault core shows that plastic creep played an important role even for generally brittle conditions. The sense of shear implied by the “ductile” structures is in agreement with the kinematic indicators on the main slip surface, indicating a consistent geometric and kinematic regime during continued movement along the Chamossaire fault.

Fig. 7 - (modified after Cardello & Mancktelow 2014). Sex Rouge south: growth fault. The Eocene deposits lie unconformably on the Wang Fm. Yellow line: palaeoescarpment. The Wang Fm passively recovers the relief by increasing sedimentation rate on the hanging wall of the no-longer active faults.

STOP 3 - (46°19'40.60"N/7°22'40.09"E) the Sex Rouge-Donin palaeo-fault

Here the Lower Cretaceous platform to ramp deposits are progressively overlapped by the Upper Cretaceous sequence, which is composed of Turonian to Campanian Amden Fm marls and the early Maastrichtian Wang Fm. This palaeo-escarpment setting is characterized by slumps and angular unconformities, related to a growth-fault structure formed during the late Cretaceous (Fig. 7). In Figure 8a, a detail of the contact between Schrattekalk fm (Urgonian limestones) and Wang Fm marls is given. This contact is marked by patchy breccias (Fig. 8b) that are interposed between the two units, with an age that probably dates back to the earliest Maastrichtian.




Fig. 8 - Details of escarpments. a) and b) Southern slope of Sex Rouge (see Fig. 7 for panoramic location). **a)** Step-like geometry of the escarpment (yellow line). The surface is silicified. **b)** Syn-sedimentary breccia preserved in the flatter parts of the escarpment underlying the onlapping deposits.




Fig. 9 - Sex Noir-Donin region. **a)** Palaeo-fault and related palaeo-escarpment crop out with **b)** slumps within the Seewen Fm. **c)** Detail of b), whose shape indicates soft-sediment deformation and gliding towards the depo-centre. **d)** Encrusted Schrattekalk fm.



STOP 4 - (46°19'51.00"N/7°22'19.22"E) half-graben structures

On the way to Stop 4, looking to the west of the valley, you can observe the Sex Noir-Donin region, which is rich in palaeographic features (Fig. 8a-c). At a height of ca. 2640 m, encrusted Urgonian limestones rich in rudists are topped by a reduced series of pelagic limestones and marls (Fig. 9d). According to our reconstructions (Fig. 10), this Stop is exactly on the southern shoulder of a half-graben, whose reduced stratigraphic thickness is due to a lateral slump of the Seewen Fm deposits, followed by hardground formation (Fig. 9d).




Fig. 10 - Cartoon summarizing the sedimentary facies and fault distribution based on profile AB in Figure 5 (Cardello & Mancktelow 2014). Numbers in brackets are **(1)** small-, **(2)**, moderate-, and **(3)** large-scale faults. Fault set **(3a)** was active in the Turonian-Santonian, whereas **(3b)** was active in the early Maastrichtian.



STOP 5 - (46°19'52.08"N/7°22'17.34"E) latest Aptian syn-sedimentary dykes

A series of NE-striking Garschella dykes and related structures, such as small scale faults and open fractures, dismember the top of the Schrattenkalk fm (Fig. 11). A graben a few decimeters across dissects the encrusted top of the Schrattenkalk fm. This is then directly covered by the Seewen Fm, which also occurs as an infilling of a few dykes oriented parallel to the Garschella related material. (Lunch break is forecast at this Stop).



Fig. 11 - Small-scale graben within Schrattenkalk fm, filled in by deposits of the Garschella Fm (Brisi member, latest Aptian) and topped by Seewen Fm (Cenomanian-Turonian).

STOP 6 - (46°19'56.94"N/7°22'9.63"E) an example of a Neogene fault

On top of the Seewen Fm and Schrattenkalk fm, an en-echelon arrangement of Neogene faults can be observed, cross-cutting the pre-existing Cretaceous syn-sedimentary faults and later fold-and-thrust structures. Some of these faults record the transition from ductile to brittle behaviour (Fig. 12).

STOP 7 - (46°19'57.98"N/7°21'57.60"E) Turonian slumps within Campanian marls

Slumps of calcareous re-sediments, which occur within the Campanian marls of the Amden Fm. (Fig. 13). Similarly to what was observed by Villars (1989) more to the east, the re-sedimented Seewen-type limestone is Turonian in age. Slumps are internally quite folded.




Fig. 12 - Examples of deformation associated with the Neogene faults. **a)** Sex Rouge eastern plateau, 2619 meters. Fault plane dips 172-63, slickenlines 240-05; **b)** Detail of Schrattekalk fm, with recrystallized and sheared cataclasite; **c)** Foliation related to the fault develops in the Seewen Fm whereas the Schrattekalk fm is affected by sheared veins and new calcite mineralization on the fault planes and finally by late fractures; **d)** Throughgoing fault and associated deformation within the Nummulitenkalk and the Wildstrubel Fm.




Fig. 13 - **a-b**) Slump deposits that are later cross-cut by Neogene veins (orange) and bedding-parallel pressure solution planes due to compaction. Black lines mark bedding and pressure solution planes. **c-d**) Close-up of tight fold produced after soft-sediment deformation.



STOP 8 - (46°20'5.15"N/7°21'37.03"E) a panoramic view of the syn-sedimentary structures

At this Stop there is the limit between the palaeo-fault and -escarpment related to the Eaux Froides syn-sedimentary fault, which accommodates nearly 0.5 km displacement and is determined clearly in the stratigraphic record Fig. 14.




Fig. 14 - **a)** Eaux Froides palaeofault structure with related palaeo-escarpment (Cardello and Mancktelow 2014). **b)** Slice of Schrattekalk preserved in the footwall of the fault. A thin veneer of Seewen Fm onlaps and subsequently covers this slice, which is then onlapped by the Amden Fm and the rest of the Tertiary succession. **c)** Fault geometry in the westernmost outcrops shows that the original structure is wavy and probably modified by later deformation.

STOP 9 - (46°20'20.50"N/7°22'25.35"E) Neogene vein sets

On the way to the hut, en-echelon veins occur associated with faults, with their number and arrangement depending on the lithology in which they are developed. Within Tertiary sandstones, kink bands can be observed. At about 17h, you should arrive at the Cabannes des Audannes (46°20'36.25"N/7°23'2.74"E, height 2500 m).



Second day: a closer look at the Cretaceous syn-sedimentary fault system

The total path length of this second day is some 7.5 km. Height difference is about 900 m downhill but with some up and down along the path, whose difficulty is medium. Some minor and quite easy out-of-path views are planned on the calcareous plateau southeast of the hut. The highest point is 2665 m. We suggest you to leave from Les Audannes hut (circa 2500 m) at 8:30 in the direction of the palaeo-escarpment, marked by the yellow dotted line in Figure 12.

STOP 10 - (46°20'57.51"N/7°23'11.25"E) the escarpment; a stratigraphic contact

At this Stop, syn-sedimentary breccia covered by a veneer of Seewen Fm and later onlapped by the Late Cretaceous to Tertiary sequence can be observed (Fig. 15).

STOP 11 - (46°21'5.56"N/7°23'15.65"E) palaeo-escarpment culmination

Palaeo-escarpment at the 2665 m culmination and a panoramic view over the Rawil pass region and the Tseuzier Lake, where the Neogene nappe-related structures are evident (Fig. 16). After this point, the way downhill to the bus stop will mostly be along the Cretaceous structures.



Fig. 15 - **a)** Detail of escarpment at Col des Eaux Froides looking to the NE. Two generations of breccia are at the contact between the Lower Cretaceous series in the footwall (left) and the Upper Cretaceous – Tertiary sequence in the hanging wall (right). A thin level of pelagic limestones (Seewen Fm) discontinuously covers the escarpment, which is later onlapped by the Amden, Wang, and the Tertiary Klimsenhorn, Wildstrubel and Globigerina Marl Fms. **b)** Detail of whitish syn-sedimentary Albian breccia.





Fig. 16 - Panoramic view of the Tseuzier Lake and valley. **a)** Thrusts crosscut the inherited Cretaceous structures (i.e., graben in the hanging wall of the main palaeofault (in violet)). During folding these structures **b)** were buttressed against the palaeo-escarpment (dotted yellow line). **c, d)** top-to-the-NW nappe-emplacement related folds and thrusts within the Kieselkalk Fm later crosscut by late normal faults (red lines; e.g., Mondraleche fault), which dip to the SSE.



Fig. 17 - Detail of Col des Eaux Froides and its palaeo-escarpment. Picture is taken from the opposite side of the valley looking to the SW. This erosive surface is defined by promontories and embayments that shape the Schrattekalk fm (on the right), which is later overlapped by the Upper Cretaceous and Tertiary succession.



STOP 12 - (46°20'48.84"N/7°23'24.04"E) palaeo-escarpment and buttressing

Similarly to what is shown in Fig 14b, the interference between the inherited Cretaceous structures and the nappe geometries is evident. At this Stop, we are right in the syncline in the hanging wall of the Eaux Froides palaeofault. Both the Wildhorn and the Ultrahelvetic nappes are folded together against the palaeo-escarpment, generating a buttressing structure (Figs. 14, 16, 17). This is further evidence that these palaeofaults preceded nappe formation.

STOP 13 - (46°20'41.86"N/7°23'27.97"E) slumps

A further Stop in the late Cretaceous slumps on the eastern shore of the Audannes Lake (Fig. 18).

STOP 14 - (46°20'37.64"N/7°23'56.62"E) a way-back trip through the Cretaceous series

A domino-like structure occurs dissecting the Lower Cretaceous series (Fig. 19). Schrattekalk fm beds are covered by a reduced and discontinuous stratigraphic series, whose thickness and facies changes according to the small- to moderate-scale faults. Similarly to what was observed the day before, the top of the Schrattekalk fm is affected by silica and Fe-rich encrustations at the top of the Lower Cretaceous series.




Fig. 18 - Detail of slump in the re-sedimented Seewen Fm type deposits within the Amden Fm. Gravity related gliding of these pelagic limestone was probably triggered by instability related to excessive steepness of the palaeo-escarpments.




Fig. 19 - (modified from Cardello & Mancktelow, 2014). The Cretaceous succession between the two major palaeofaults is formed by a rollover anticline, broken by a syn-sedimentary domino-like structure, later crosscut by the thrust trajectories. Red thin lines are late Neogene faults.



STOP 15 - (46°20'7.12"N/7°23'7.79"E) palaeo-geographic Maastrichtian evidences

A close look at a step of the Sex Rouges-Donin palaeofault that was completely sealed during the sedimentation of the Wang Fm in the early Maastrichtian (Fig. 20).

After this Stop, the bus stop can be reached to take the way back to Sion (Fig. 1, Stop 16 - 46°19'57.31"N/7°26'14.63"E).




Fig. 20 - **a)** Moderate scale paleofault sealed during the late Cretaceous. **b)** Detail of the Upper Cretaceous succession. Away from the edge of the normal fault, re-sediments are thicker and lenses of calcarenites occur. **c)** A patch of pebbly mudstone of Seewen Fm type limestone along the palaeo-escarpment surface.



The Wildhorn Nappe structure from Cretaceous to Neogene time

*8th - 9th August 2015
summer excursion organized by
Giovanni Luca Cardello & Neil Mancktelow*



ACKNOWLEDGEMENTS

The research presented on this field excursion was supported by Swiss National Science Foundation projects 200020-121578 and 200020-137510.



References

- Burkhard M. (1988) - L'Helvétique de la bordure occidentale du massif de l'Aar (évolution tectonique et métamorphique). *Eclogae Geologicae Helveticae*, 81, 63-114.
- Burkhard M. (1993) - Calcite twins, their geometry, appearance and significance as stress-strain markers and indicators of tectonic regime: a review. *Journal of Structural Geology*, 15(3), 351-368.
- Burkhard M. & Kerrich R. (1988) - Fluid regimes in the deformation of the Helvetic Nappes, Switzerland, as inferred from stable isotope data. *Contributions to Mineralogy and Petrology*, 99, 416-429.
- Campani M., Mancktelow N. & Courrioux G. (2014) - The 3D interplay between folding and faulting in a syn-orogenic extensional system: the Simplon Fault Zone in the Central Alps (Switzerland and Italy). *Swiss Journal of Geosciences*, 107(2-3), 251-271.
- Cardello G.L. (2013) - The Rawil depression: its structural history from Cretaceous to Neogene. Ph.D. dissertation, Switzerland. ETH Zurich. doi:10.3929/ethz-a-010016215
- Cardello G.L., & Tessei T. (2013) - Transtensive faulting in carbonates at different crustal levels: examples from SW Helvetic and Central Apennines. *Rendiconti online della Società Geologica Italiana*, 29, 20-23.
- Cardello G.L. & Doglioni C. (2015) - From Mesozoic rifting to Apennine orogeny: The Gran Sasso range (Italy). *Gondwana Research*, 27(4), 1307-1334.
- Cardello G.L. & Mancktelow N.S. (2014) - Cretaceous syn-sedimentary faulting in the Wildhorn Nappe (SW Switzerland). *Swiss Journal of Geosciences*, 107(2-3), 223-250.
- Cardello G.L. & Mancktelow N.S. (2015) - Veining and post-nappe transtensional faulting in the SW Helvetic Alps (Switzerland). *Swiss Journal of Geosciences*, 108(2-3), 379-400.
- Cardello G.L., Almqvist S.B, Hirt A.M. & Mancktelow N.S. (2015) - Determining the timing of formation of the Rawil Depression in the Helvetic Alps with the use of paleomagnetic and structural methods. *Geological Society Special Publication*, SP 425(4). doi:10.1144/SP425.10
- Casey M., Dietrich D. & Ramsay J.G. (1983) - Methods for determining deformation history for chocolate tablet boudinage with fibrous crystals.
- Crespo-Blanc A., Masson H., Sharp Z., Cosca M. & Hunziker J. (1995) - A stable and $^{40}\text{Ar}/^{39}\text{Ar}$ isotope study of a major thrust in the Helvetic nappes (Swiss Alps): Evidence for fluid flow and constraints on nappe kinematics. *Geological Society of America Bulletin*, October 1995, 107(10), 1129-1144.
- Diehl T., Deichmann N., Clinton J., Husen S., Kraft T., Plenkers K., Guilhelm A., Behr Y., Cauzzi C., Kästi P., Haslinger F., Fäh D., Michel C., Wiemer S. & Woessner J. (2013) - Earthquakes in Switzerland and surrounding regions during 2012. *Swiss Journal of Geosciences*, 106(3), 543-558.
- Dietrich D., McKenzie J.A. & Song H. (1983) - Origin of calcite in syntectonic veins as determined from carbon-isotope ratios. *Geology*, 11, 547-551.
- Durney D.W. & Ramsay J.G. (1973) - Incremental strains measured by syntectonic crystal growths. In: K.A. de Jong & R. Scholten, *Gravity and tectonics*, 67-96, Wiley.

- Egli D. & Mancktelow N. (2013) - The structural history of the Mont Blanc massif with regard to models for its recent exhumation. *Swiss Journal of Geosciences*, 106, 469-489.
- Franck P., Wagner J.J., Escher A. & Pavoni N. (1984) - Evolution des contraintes tectoniques et sismicité dans la région du col du Sanetsch, Alpes valaisannes helvétiques. *Eclogae Geologicae Helvetiae*, 77(2), 383-393.
- Frischknecht C., Rosset P. & Wagner J.J. (2005) - Toward seismic microzonation—2-D modeling and ambient seismic noise measurements: The case of an embanked, deep alpine valley. *Earthquake spectra*, 21(3), 635-651.
- Gasser D., & Mancktelow N.S. (2010) - Brittle faulting in the Rawil depression: field observations from the Rezli fault zones, Helvetic nappes, Western Switzerland. *Swiss Journal of Geosciences*, 103, 15-32.
- Glotzbach C., Van Der Beek P.A. & Spiegel C. (2011) - Episodic exhumation and relief growth in the Mont Blanc massif, Western Alps from numerical modelling of thermochronology data. *Earth and Planetary Science Letters*, 304(3), 417-430.
- Gubler E., Kahle H.G., Klingele E., Müller S., & Olivier R. (1981) - Recent crustal movements in Switzerland and their geophysical interpretation. *Tectonophysics*, 38, 297-315.
- Günzler-Seiffert H. (1941) - Persistente Brücke im Jura der Wildhorn-Decke des Berner Oberlandes. *Eclogae Geologicae Helvetiae*, 34/2. 164 -172.
- Heim A. (1920) - *Geologie der Schweiz*. B. Tauchnitz Verlag GmbH, Leipzig.
- Hänni R. & Pfiffner O.A. (2001) - Evolution and internal structure of the Helvetic nappes in the Bernese Oberland. *Eclogae Geologicae Helvetiae*, 94, 161-171.
- Hoffmann B.A., Helfer M., Diamond L.W., Villa I.M., Frei R. & Eikenberg J. (2004) - Topography-driven hydrothermal breccia mineralization of Pliocene age at Grimsel Pass, Aar massif, Central Swiss Alps. *Schweizerische Mineralogische und Petrographische Mitteilungen*, 84, 271-302.
- Jeanbourquin P. (1994) - Early deformation of Ultrahelvetique mélanges in the Helvetic nappes (Western Swiss Alps). *Journal of Structural Geology*, 16/10, 1367-1383.
- Kastrup U., Zoback M.L., Deichmann N., Evans K.F., Giardini D. & Michael A.J. (2004) - Stress field variations in the Swiss Alps and the northern Alpine foreland derived from inversion of fault plane solutions. *Journal of Geophysical Research*, 109, B01402, doi:10.1029/2003JB002550.
- Kempf O. & Pfiffner O.A. (2004) - Early Tertiary evolution of the North Alpine Foreland Basin of the Swiss Alps and adjoining areas. *Basin Research*, 16(4), 549-567.
- Kirschner D.L., Masson H. & Cosca M.A. (2003) - An $^{40}\text{Ar}/^{39}\text{Ar}$, Rb/Sr, and stable isotope study of micas in low-grade fold-and-thrust belt: an example from the Swiss Helvetic Alps. *Contributions to Mineralogy and Petrology* 145, 460-480.
- Lugeon M. (1914-1918) - Les Hautes Alpes calcaires entre la Lizerne et la Kander: (Wildhorn, Wildstrubel, Balmhorn et Torrenthorn): (explication de la carte spéciale no. 60). 3 Fascicule. Matériaux pour la carte géologique de la Suisse. Nouvelle série 60.
- Maurer H.R., Burkhard M., Deichmann N. & Green A.G. (1997) - Active tectonism in the central Alps: contrasting stress regimes north and south of the Rhone Valley. *Terra Nova*, 9, 91-94.
- Mohn G., Manatschal G., Müntener O., Beltrando M. & Masini E. (2010) - Unravelling the interaction between tectonic and sedimentary processes during lithospheric thinning in the Alpine Tethys margins. *International Journal of Earth Sciences*, 99 (Suppl. 1), 75-101.

- Pavoni N. (1980) - Comparison of focal mechanisms of earthquakes and faulting in the Helvetic zone of the Central Valais, Swiss Alps. *Eclogae Geologicae Helveticae*, 73, 551-558.
- Pfiffner O.A. (2009) - *Geologie der Alpen*. Bern/Stuttgart/Wien: Haupt Verlag, 359 pp.
- Ramsay J.G., Casey M., Dietrich D., Kligfield R., Mancktelow N., Schmid S., Siddans A.W.B. & Pfiffner A.O. (1981) - Strain features of the Helvetic nappes of Switzerland. In: *The Geological Society of America, 94th annual meeting 13. Abstracts with Programs - Geological Society of America*, 535.
- Ramsay J.G., Casey M. & Kligfield R. (1983) - Role of shear in development of the Helvetic fold-thrust belt of Switzerland. *Geology*, 11, 439-442.
- Steck A., Bigioggero B., Dal Piaz G.V., Escher A., Martinotti G. & Masson H. (1999) - Carte géologique des Alpes de Suisse occidentale, 1:100,000, Carte géologique spéciale N°123. Berne: Service Hydrologique et Géologique National.
- Sternai P., Herman F., Champagnac J.D., Fox M., Salcher B. & Willett S.D. (2012) - Pre-glacial topography of the European Alps. *Geology*, 40(12), 1067-1070.
- Sue C., Delacou B., Champagnac J.-D., Allanic C., Tricart P. & Burkhard M. (2007) - Extensional neotectonics around the bend of the Western/Central Alps: an overview. *Int. J. Earth Sci. (Geol Rundsch)*, 96(6), pp. 1101-1029. doi:10.1007/s00531-007-0181-3
- Trümpy R. (1960) - Paleotectonic evolution of the central and western Alps. *Geological Society of America Bulletin*, 71, 843-908.
- Ustaszewski M., McClymont A., Herwegh M., Preusser F. & Pfiffner O.A. (2007) - Unravelling the evolution of an Alpine to post-glacially active fault in the Swiss Alps. *Journal of Structural Geology*, 29-12, 1943-1959. doi:10.1016/j.jsg.2007.09.006
- Vernon A.J., Van Der Beek P.A., Sinclair H.D. & Rahn M.K. (2008) - Increase in late Neogene denudation of the European Alps confirmed by analysis of a fission-track thermochronology database. *Earth and Planetary Science Letters*, 270(3), 316-329.
- Villars F. (1989) - Mégablocs resédimentés dans les Marnes d'Amden de la Plaine Morte (nappe helvétique du Wildhorn, Suisse): tectonique synsédimentaire à la fin du Santonien. *Schweizerische Mineralogische und Petrographische Mitteilungen*, 69, 167-172.